



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2009

---

## **Long-term effects of early parasite exposure on song duration and singing strategy in great tits**

Bischoff, Linda ; Tschirren, Barbara ; Richner, Heinz

**Abstract:** Song is a sexually selected trait in many bird species and has been suggested to function as a signal of a male's health and parasite resistance. Here we present an experimental field study on the long-term effects of parasite exposure early in life on adult bird song. We exposed nestling great tits (*Parus major*) to ectoparasitic hen fleas (*Ceratophyllus gallinae*) and on their recruitment into the local breeding population assessed the response to a playback of a challenging male in their breeding territory. We show, to our knowledge for the first time in a wild bird population, that parasite exposure early in life affects bird song: song duration of males that were exposed to parasites early in life was reduced by 32% compared with males that grew up in a flea-free environment. Early parasite exposure also significantly reduced the degree of song overlap with the playback, which has been shown to correlate with social status. There was no effect of early parasite exposure on the number of different song types sung or on the latency until the males started the vocal response to the playback. These results suggest that mates or rivals can use song duration and song overlap as proxies for an individual's exposure to parasites early in life. It thereby highlights both the importance of parasites in maintaining honesty of sexually selected traits and the costs of parasitism in terms of reduced attractiveness and competitiveness.

DOI: <https://doi.org/10.1093/beheco/arp012>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-76104>

Journal Article

Accepted Version

Originally published at:

Bischoff, Linda; Tschirren, Barbara; Richner, Heinz (2009). Long-term effects of early parasite exposure on song duration and singing strategy in great tits. *Behavioural Ecology*, 20:265-270.

DOI: <https://doi.org/10.1093/beheco/arp012>

**Longterm effects of parasite exposure early in life on song duration and singing strategy – a case study in great tits**

Linda L. Bischoff<sup>a\*</sup>, Barbara Tschirren<sup>b</sup> & Heinz Richner<sup>a</sup>

<sup>a</sup>Evolutionary Ecology Lab, Ecology and Evolution Institute, University of Bern, Baltzerstrasse 6, 3012 Bern, Switzerland

<sup>b</sup>Centre for the Integrative Study of Animal Behaviour, Macquarie University, Sydney, NSW 2109, Australia

\* author for correspondence: linda.bischoff@esh.unibe.ch

Short title: Parasites affect bird song

Key words: bird song; ectoparasites; developmental stress hypothesis; honest signaling; *Parus major*; sexual selection

Word count: 5621

## ABSTRACT

Song is a sexually selected trait in many bird species, and has been suggested to function as a signal for a male's health and parasite resistance. Here we present an experimental field study on the long-term effects of parasite exposure early in life on adult bird song. We exposed nestling great tits (*Parus major*) to ectoparasitic hen fleas (*Ceratophyllus gallinae*), and upon their recruitment into the local breeding population assessed the response to a playback of a challenging male in their breeding territory.

The duration of the singing response of breeding males that had undergone post-natal exposure to parasites was reduced by 32% compared to males that grew up in a parasite-free environment, and also significantly reduced the degree of song overlap with the playback, which is thought to correlate with social status. There was no effect of early parasite exposure on the number of different song types sung within 20 minutes, or on the latency until the males started the vocal response to the playback. The results thus suggest, to our knowledge for the first time in an experimental field study, that mates or rivals can use song duration and song overlap as proxies for an individual's exposure to parasites early in life.

## 1. INTRODUCTION

The song of male birds often has a dual signalling function in both female attraction and repulsion of competing males (Darwin 1871; Catchpole 1982; Searcy and Andersson 1986; Catchpole and Slater 1995). Singing is energetically costly for males (Brackenbury 1979; Lambrechts and Dhondt 1988; Vehrencamp et al. 1989; Grafen 1990; Eberhardt 1994), implying that higher quality males can afford to display more costly song types, song structures or song rates. Numerous studies have shown that a male's present condition influences song rate and song repertoire (e.g., Buchanan et al. 1999; Berg et al. 2005). More recently, it has been suggested that a male's song might also be affected by environmental conditions experienced early in life. Indeed, several studies (Nowicki et al. 2002; Buchanan et al. 2003) have shown that nutritional stress during early development can affect song development (i.e., the nutritional stress hypothesis, Nowicki et al. (1998)). This hypothesis has been expanded to other stress factors, for example early parasite exposure or increased corticosterone levels, that might have long-lasting effects on song characteristics in adulthood (i.e., the developmental stress hypothesis, Spencer et al. (2003); Buchanan et al. (2004); Nowicki & Searcy, (2004); but see Gil (2006)).

Song has thereby the potential to function as a multiple signal of both a male's past growth and development (Buchanan et al. 2003; Spencer et al. 2003; Spencer et al. 2005a), and / or of his current condition (Buchanan et al. 1999). By assessing song rates and song structures during mate choice, females might thus not only gain honest information about a male's current status, but potentially also on past environmental conditions experienced by the male early in life (Catchpole 1996; Buchanan 2000; Buchanan and Catchpole 2000; Doutrelant et al. 2000).

63           Honest information about male quality is not only relevant for female mate  
64 choice, but also for male-male competition. Males often eavesdrop on the neighbour's  
65 song to get honest information on the motivation, condition or health status of poten-  
66 tial contrahents (Peake et al. 2005). Two important aspects of intra-sexual vocal sig-  
67 nalling are song matching, where an individual repeats the song pattern of the com-  
68 petitor (Krebs et al. 1981; McGregor et al. 1992; Nielsen and Vehrencamp 1995; Bee-  
69 cher et al. 2000; Vehrencamp 2001), and song overlapping, where an individual, de-  
70 pending on its relative dominance status, times its song relative to the opponent's  
71 song with various degrees of overlap (Dabelsteen et al. 1997).

72           To date, studies testing the effects of early environmental conditions on song  
73 development have been performed mainly under lab or aviary conditions (Buchanan  
74 et al. 2003; Spencer et al. 2004), while in the wild evidence for effects of stress expe-  
75 rienced early in life on song development remains scarce. Here we investigated if  
76 early developmental stress, resulting from exposure to a common ectoparasite during  
77 the nestling period, can affect song traits and song overlap of males under natural  
78 conditions. We thereto experimentally infested nestling great tits (*Parus major*) after  
79 hatching with hen fleas (*Ceratophyllus gallinae*) and recorded their song after re-  
80 cruitment into the local breeding population the following years. Hen fleas are com-  
81 mon ectoparasites of great tits. They live in the nest material of their hosts and suck  
82 blood from the nestlings (Tripet and Richner 1997b). If early developmental condi-  
83 tions have long-term consequences for adult song we predict a shorter duration of  
84 singing and / or a smaller number of different song types sung within the recording  
85 period in males that were exposed to hen fleas early in life. Furthermore, we predict  
86 that males raised in a parasite-free nest are more competitive and thus show a higher  
87 degree of song overlap with a competing male.

88

## 89 **2. MATERIAL AND METHODS**

### 90 *a) Manipulation of early parasite exposure*

91 The study was performed in a population of great tits breeding in nest-boxes in a for-  
92 est close to Bern, Switzerland („Forst“, 46°54'N 7°17'E / 46°57'N 7°21'E). During  
93 the breeding seasons of 1998 – 2004 we manipulated the load of hen fleas in the nests.  
94 First, we microwaved the nesting material of all nests two days after hatching of the  
95 nestlings. This procedure kills all nest-based ectoparasites naturally present in the  
96 nests (Richner et al. 1993; Tripet and Richner 1997a). Thereafter, half of the nests  
97 were experimentally infested with 40 female and 20 male fleas originating from old  
98 nests collected within the same forest before the breeding season, while the other  
99 nests remained flea-free. Nests were randomly assigned to one of the two treatment  
100 groups. All nestlings were ringed with individually numbered aluminium rings when  
101 9 days old. Fifteen days post-hatching, towards the end of the nestling stage, the nest-  
102 lings' body mass and metatarsus length were measured.

103 Twenty-two males (13 from experimentally infested nests and 9 from nests  
104 without parasites) were recaptured as breeders in 2005 in the local population. All  
105 recruits originated from different broods (i.e., they were not siblings). One of these  
106 males was born and manipulated in 1998 (no fleas), three males were born and ma-  
107 nipulated in 2002 (two with fleas and one without fleas), and the rest of the males  
108 were born and manipulated in 2004. In summary, our sample consisted of 21 males,  
109 17 of them were 1<sup>st</sup> year breeders and four of them older. Of the four older birds, two  
110 had been experimentally infested as nestling and two came from uninfested controls.

111

### 112 *b) Recording and analysis of song*

To create the song stimuli, we recorded the songs of five males from outside of our study area (“Birchiwald”, Bremgarten b. Bern, 46°58’N 7°25’E, Switzerland, distance to study area ~10 km) from a distance of approximately 10 – 15 meters during 45 minutes each. The songs were recorded with a condenser shot gun microphone (Georg Neumann, KMR 82 i) via a preamplifier (SHURE, FP23) to the program Sound Studio 2.1 (Kwok 2003) on a laptop. We used these digitised songs to create five different song stimuli of 20 minutes duration in the programs Sound Studio 2.1 (Kwok 2003) and Raven 1.2.1 (Charif et al. 2004) (see details below).

The song of great tits has a typical structure (Figure 1): a combination of 1 – 5 notes (Krebs 1976; McGregor and Krebs 1982b) is called a phrase (about 0.2 – 0.5 sec; Figure 1). Phrases are usually repeated and form a strophe (McGregor and Krebs 1982b; Lambrechts and Dhondt 1986). Strophes are typically separated by an interstrophe pause before they are repeated again. Repeated strophes based on the same kind of phrase are called a song type (sometimes also called song unit, Lambrechts and Dhondt (1988)). A male great tit has a song repertoire of 1 to 9 distinct song types (McGregor et al. 1981; McGregor and Krebs 1982a; Otter et al. 1999). Females are attracted to males with larger song repertoires (Baker et al. 1986). Generally, great tits sing strophes of the same song type for several minutes before switching to another song type (Krebs 1976).

Some song types are much more common than others. The most common song type in our study population (Figure 1) also occurred in all recordings of the five birds used to create the different song stimuli. Single phrases of these most common song type were extracted and multiplied six times using Sound Studio 2.1 (Kwok 2003) to create a stimulus strophe of 4 seconds followed by 3 seconds of silence (i.e., the interstrophe pause, Figure 1). The duration of strophe and pause was based on a pre-

liminary study and together represents an average male's song unit in the study population (L.B. unpublished data). The song unit (i.e., the strophe and pause) was then repeated to create 1-minute stimulus sequences. Each stimulus contained the song of one male only. This approach allowed us to standardize the stimulus structure, while still covering a range of natural variation in song between different males. The amplitude of the five song stimuli were very similar, that is  $83 \pm 2$  dB, measured at a distance of 1m in front of the loudspeaker using a sound pressure level meter (Votcraft Plus, SL-300).

We recorded the males' response to a playback during the breeding season 2005. All recordings were performed between 6am – 1pm on the day the clutch contained the fourth egg. We played back a non-interactive 20-minute song stimulus in the male's breeding territory using an iPod mini with the full uncompressed WAVE/AIFF format setting and a loudspeaker (YAMAHA, AA5), which was placed one meter away from the nest box on the ground. The amplitude of the stimulus songs was standardized by using fixed settings on both the playback machine and the speaker.

For the song recordings of the target males, we first played a 5-minute song stimulus sequence (as described above) in each male's territory, followed by 30 seconds of silence. Then, we played six times a 2-minute song stimulus followed by a 30 seconds of silence to make up for a total of 20 minutes of playback in each male's territory. Thus each target male was exposed to the song of one male used as a stimulus selected at random from the five available stimuli (systematic randomization to assure that all stimuli were used equally often). All stimuli were played back to males originating from both parasite-infested and parasite-free nests respectively (stimulus 1: without fleas  $n = 3$ , with fleas  $n = 2$ ; stimulus 2: without fleas  $n = 1$ , with fleas  $n =$



3; stimulus 3: without fleas  $n = 2$ , with fleas  $n = 2$ ; stimulus 4: without fleas  $n = 1$ , with fleas  $n = 3$ ; stimulus 5: without fleas  $n = 2$ , with fleas  $n = 3$ ), and stimulus number (1-5) was included as a random effect in the statistical analyses to control for variation in the males' response due to variation among stimuli.

The song of the focus males was recorded during the entire 20-minute trial period using a condenser shot gun microphone (Georg Neumann, KMR 82 i) via a pre-amplifier (SHURE, FP23) to the program Sound Studio 2.1 (Kwok 2003) on a laptop. The spectrograms were analysed blindly with respect to the parasite treatment of the males (at the time they were nestlings) using the program Raven 1.2.1 (Charif et al. 2004). We measured the summed duration of all song units (see Figure 1 and above for the definition of a song unit, i.e. the combined duration of strophe plus interstrophe pause) during the 20 minute recording (thereafter termed song duration), the number of different song types sung within these 20 minutes, the latency until the focus male responded to the playback, the summed duration of all strophe lengths during the 20 minutes (termed after stroph length), and the summed duration of all interstrophe pauses during the 20 minutes (termed after interstrophe pause length). We also assessed the number of times that a target male sang on top of the stimulus in order to calculate song overlap as follows: only cases where the target bird responded *after* the stimulus strophe (for definition of song strophe see Figure 1) had started were included, but all cases where the target male started *before* the stimulus strophe were discarded from this analysis since in this cases song overlap may have arisen just by chance. Percentage of song overlap was then calculated as the number of overlapping strophes / number of total strophes \*100.

### *c) Clutch size and breeder morphology*

Clutch size was recorded at the start of incubation. Breeding parents were captured 12 days after hatching of their brood using a door trap inside the nest box, and their metatarsus length and body mass were measured. Their body condition was calculated as the residuals of a regression of body mass on tarsus length. The change in body mass (Repeated measures ANOVA:  $F_{1, 20} = 1.957$ ,  $P = 0.177$ ) and metatarsus length (Repeated measures ANOVA:  $F_{1, 20} = 2.870$ ,  $P = 0.106$ ) between fledging and adulthood did not differ significantly between males that had been parasite-exposed or unexposed as nestlings.

#### *d) Statistical analyses*

We used general linear models to analyse the effect of early flea exposure on the song duration, the number of different song types sung within these 20 minutes, the latency until the male started their vocal response to the playback, the strophe length, the interstrophe pause length, and the percentage of song overlap. Song stimulus type (1 – 5) was included as a random factor in all statistical analyses to control for differences in a male's vocal response due to variation among playback stimuli. Residuals of the models were tested for normality using Shapiro-Wilk tests and homoscedasticity using Bartlett tests to ensure that the assumptions for parametric testing were fulfilled. All tests were two-tailed with a significance level of 0.05. Means  $\pm$  1 SE are presented in the results and the figures. Statistical analyses were performed in the programs R 2.6 (R Development Core Team 2004) and JMP IN 5.1 (Sall and Lehmann 1996).

### **3. RESULTS**

#### *a) Effects of experimental flea exposure on song*

Males raised in a parasite-infested nest responded vocally for a significantly shorter period ( $364 \pm 55$  sec,  $n = 13$ ) during the 20 minute playback trials than males brought up in a parasite-free nest ( $536 \pm 39$  sec,  $n = 9$ ;  $F_{1, 16} = 5.396$ ,  $P = 0.034$ ). This longer song duration of parasite-free birds was due to a tendency for longer strophe length ( $F_{1, 16} = 4.179$ ,  $P = 0.057$ ) as well as longer interstrophe pauses ( $F_{1, 16} = 8.595$ ,  $P = 0.009$ ). Males raised in a parasite-free nest also overlapped their song significantly more with the playback ( $62.9\% \pm 4.4$ ) than males from an infested nest ( $48.3\% \pm 2.4$ ;  $F_{1, 16} = 10.209$ ,  $P = 0.006$ ). This effect became even more significant ( $F_{1, 15} = 21.972$ ,  $P < 0.001$ ) when including song duration ( $F_{1, 15} = 8.634$ ,  $P = 0.010$ ) as a covariate into the analysis of song overlap, showing that early flea exposure influenced song overlap and song duration independently (Figure 2).

Early flea exposure did not significantly influence the number of different song types sung within these 20 minutes (with fleas:  $3.3 \pm 0.6$ , without fleas:  $3.0 \pm 0.4$ ;  $F_{1, 16} = 0.171$ ,  $P = 0.684$ ), or the latency until the males started their vocal response to the playback (with fleas:  $170.1 \pm 63.9$  sec, without fleas:  $148.2 \pm 98.3$  sec;  $F_{1, 16} = 0.038$ ,  $P = 0.847$ ).

#### ***b) Correlates of song***

Song duration during the playback trials was not significantly influenced by a male's body condition as an adult ( $F_{1, 16} = 0.137$ ,  $P = 0.716$ ) or as a nestling ( $F_{1, 16} = 0.004$ ,  $P = 0.948$ ), and it was independent of male age ( $F_{1, 16} = 0.003$ ,  $P = 0.954$ ), or the date ( $F_{1, 16} = 0.408$ ,  $P = 0.532$ ) or time of day of the song recording ( $F_{1, 16} = 0.475$ ,  $P = 0.501$ ).

Similarly, song overlap was not significantly influenced by a male's nestling body condition ( $F_{1, 16} = 0.094$ ,  $P = 0.763$ ), male age ( $F_{1, 16} = 0.848$ ,  $P = 0.371$ ), or the

time of day of song recording ( $F_{1, 16} = 0.147$ ,  $P = 0.706$ ). We found a tendency for males in better adult body condition to overlap their song more with the playback ( $F_{1, 16} = 3.846$ ,  $P = 0.068$ ), and also a decrease of song overlap with the ongoing breeding season ( $F_{1, 16} = 4.275$ ,  $P = 0.055$ ). The effect of the flea treatment on song overlap remained significant ( $F_{1, 14} = 8.274$ ,  $P = 0.012$ ) when including adult condition and date of song recording as covariates into the statistical model.

Females laid larger clutches when mated to a male with a longer song duration ( $F_{1, 16} = 5.138$ ,  $P = 0.038$ , Figure 3). No other aspect of song was significantly correlated with clutch size (all  $P > 0.160$ ) or male adult condition ( $P = 0.343$ ). There was no significant correlation between female and male condition ( $F_{1, 19} = 0.387$ ,  $P = 0.541$ ), indicating that there was no assortative mating for body condition.

#### 4. DISCUSSION

Several studies have shown that the brain structures associated with song learning are costly to produce and thus vulnerable to stress (e.g., Catchpole 1996; Nowicki et al. 1998; but see Gil et al. 2006). In swamp sparrows (*Melospiza georgina*), for example, nutritional limitation during growth had a negative effect on the development of the song nuclei in the brain (Nowicki et al. 2002). Similarly, nestling development was a predictor of adult song repertoire size in great reed warblers (Nowicki et al. 2000). Furthermore, male european starlings (*Sturnus vulgaris*) that experienced an unpredictable food supply early in life produced a smaller repertoire of song phrases than males with a constant food supply (Spencer et al. 2004).

Beside these well documented effects of nutritional stress on song development, a recent study by Spencer et al. (2005a) demonstrated that an infection with blood parasites early in life can similarly affect song development. Captive male ca-

naries (*Serinus canaria*) that were infected with malaria parasites (*Plasmodium relictum*) during development produced simpler songs (number of different syllables) as adults and had a significantly smaller volume of the HVC nucleus (i.e., the “high vocal centre” in the brain) compared to uninfected males. Our study demonstrates that similar effects of early parasite exposure on song development occur in the wild as well: Male great tits that had been exposed to ectoparasitic hen fleas early in life sang for a shorter period during the playback trials than unexposed males. This is in agreement with the hypothesis that song development is costly (Eberhardt 1994; Gaunt et al. 1996), and it indicates that parasites might mediate the trade-off between investing limited resources in the development of song-related brain structures and parasite defence respectively. It thereby highlights the importance of parasites in ensuring honesty of sexually selected traits (Hamilton and Zuk 1982).

Interestingly, the effect of early parasite exposure on song duration were independent of adult or nestling condition. Furthermore, the number of different song types sung within the recording period and the latency until the male started their vocal response to the playback were not affected by the infestation experiment. This is surprising because Spencer et al. (2004, 2005a) have found that nutritional stress affects song repertoire size in passerines. It suggests that food restriction and parasite exposure affect song development in different ways, and it shows that song does not only contain multiple information about a male’s condition over different time-scales (see also Scheuber et al. 2003a, 2003b), but also specific information about the specific type of stressor a male was exposed to early in life.

We found a positive correlation between a male’s song duration and his partner’s clutch size, which indicates that the duration of singing is a sexually selected trait in our study population (see also Houtman 1992; Collins et al. 1994; Gil and Gahr 2002;

Garamszegi and Møller 2004; Spencer et al. 2005b). This effect is unlikely due to assortative mating because male and female condition were not significantly correlated. It thus demonstrates the benefits of a high song output for a male's fitness and reveals the cost of parasitism for males in terms of reduced attractiveness.

Beside the effects of early ectoparasite exposure on the duration of singing, which might be especially important in inter-sexual selection (see above), we also found that parasite-exposed males overlapped the song of a challenger (i.e., the playback song stimulus) less than unexposed males. Several studies have demonstrated that song overlapping is important in male-male competition as a signal of aggressiveness (Todt 1981; Brindley 1991; McGregor et al. 1992; Dabelsteen et al. 1996; Dabelsteen et al. 1997; Naguib et al. 1999; Langemann et al. 2000; Mennill and Ratcliffe 2004b), and dominance (Hyman et al. 2004; Mennill and Ratcliffe 2004b, 2004a). Males that overlap their opponent's song in male-male competition may therefore be more likely to find a mating partner or to gain access to higher quality territories, and might thereby achieve a higher reproductive success. In support of this hypothesis, Kunc et al. (2006) found that the subsequent pairing success of male nightingales (*Luscinia megarhynchos*) was related to the frequency of song overlap in male-male interactions during the time of mate attraction. Also, in a dual-speaker choice experiment female little blue penguins (*Eudyptula minor*) preferred the song of males with a higher degree of song overlap (Miyazaki and Waas 2002). Furthermore, in an experiment with alternating and overlapping playback, corn bunting males (*Miliaria calandra*) approached the loudspeaker more quickly and spent more time close to the loudspeaker when the playback was alternated rather than when it was overlapping with their own song (Osiejuk et al. 2007), indicating that competitors are less likely to challenge overlapping singers (Todt and Naguib 2000; Osiejuk et al. 2004).

Together, these studies suggests that parasite exposure early in life does not only affect a male's attractiveness, but also his territory defence behaviour and competitiveness in intra-sexual encounters by altering its singing strategy.

To summarise, we experimentally show, to our knowledge for the first time in a wild bird population, that parasite exposure early in life affects song duration and the timing of singing of adult great tits. It demonstrates that song development is costly and highlights both the importance of parasites in maintaining honesty of sexually selected traits and the costs of parasitism in terms of reduced attractiveness and competitiveness.

## **ACKNOWLEDGEMENTS**

We thank Martin C. Fischer for the assistance with the song analysis and for comments during preparation of the manuscript, Verena Saladin for field assistance and comments on the manuscript, and Katharina Gallizzi for comments on statistical questions. The experiments were conducted under a licence of the Ethical Committee of the Office of Agriculture and Nature of the Canton Bern, Switzerland. The study was financially supported by a Swiss National Science Foundation research grant (no 3100A0-102017 to Heinz Richner). Barbara Tschirren was financially supported by an Australian Research Council APD Fellowship (no DP0879313).

332   **REFERENCES**

- 333   Baker MC, Bjerke TK, Lampe H, Espmark Y. 1986. Sexual-responses of female great  
334       tits to variation in size of males song repertoires. *Am Nat.* 128:491-498.
- 335   Beecher MD, Campbell SE, Burt JM, Hill CE, Nordby JC. 2000. Song-type matching  
336       between neighbouring song sparrows. *Anim Behav.* 59:21-27.
- 337   Berg ML, Beintema NH, Welbergen JA, Komdeur J. 2005. Singing as a handicap: the  
338       effects of food availability and weather on song output in the australian reed  
339       warbler *Acrocephalus australis*. *J Avian Biol.* 36:102-109.
- 340   Brackenbury JH. 1979. Power capabilities of the avian sound-producing system. *J*  
341       *Exp Biol.* 78:163-166.
- 342   Brindley EL. 1991. Response of european robins to playback of song - neighbour  
343       recognition and overlapping. *Anim Behav.* 41:503-512.
- 344   Buchanan KL. 2000. Stress and the evolution of condition-dependent signals. *Trends*  
345       *Ecol Evol.* 15:156-160.
- 346   Buchanan KL, Catchpole CK. 2000. Song as an indicator of male parental effort in  
347       the sedge warbler. *Proc R Soc Lond B Biol Sci.* 267:321-326.
- 348   Buchanan KL, Catchpole CK, Lewis JW, Lodge A. 1999. Song as an indicator of par-  
349       asitism in the sedge warbler. *Anim Behav.* 57:307-314.
- 350   Buchanan KL, Leitner S, Spencer KA, Goldsmith AR, Catchpole CK. 2004. Devel-  
351       opmental stress selectively affects the song control nucleus HVC in the zebra  
352       finch. *Proc R Soc Lond B Biol Sci.* 271:2381-2386.
- 353   Buchanan KL, Spencer KA, Goldsmith AR, Catchpole CK. 2003. Song as an honest  
354       signal of past developmental stress in the european starling (*Sturnus vulgaris*).  
355       *Proc R Soc Lond B Biol Sci.* 270:1149-1156.



356 Catchpole CK. 1982. The evolution of bird sounds in relation to mating and spacing  
 357 behaviour. In: Kroodsma DE, Miller EH, editors. Acoustic Communication in  
 358 Birds. New York: Academic Press. p. 297-319.  
 359 Catchpole CK. 1996. Song and female choice: good genes and big brains? Trends  
 360 Ecol Evol. 11:358-360.  
 361 Catchpole CK, Slater PJB. 1995. Bird song. Biological themes and variations. Cam-  
 362 bridge: Cambridge University Press.  
 363 Charif RA, Clark CW, Firststrup KM. 2004. Raven 1.2 User's Manual. New York:  
 364 Cornell Laboratory of Ornithology.  
 365 Collins SA, Hubbard C, Houtman AM. 1994. Female mate choice in the zebra finch -  
 366 the effect of male beak color and male song. Behav Ecol Sociobiol. 35:21-25.  
 367 Dabelsteen T, McGregor PK, Holland J, Tobias JA, Pedersen SB. 1997. The signal  
 368 function of overlapping singing in male robins. Anim Behav. 53:249-256.  
 369 Dabelsteen T, McGregor PK, Shepherd M, Whittaker X, Pedersen SB. 1996. Is the  
 370 signal value of overlapping different from that of alternating during matched  
 371 singing in great Tits? J Avian Biol. 27:189-194.  
 372 Darwin C. 1871. The Descent of Man and Selection in Relation to Sex. London: Mur-  
 373 ray.  
 374 Doutrelant C, Blondel J, Perret P, Lambrechts MM. 2000. Relationship between song  
 375 repertoire size and male quality in blue tits. J Avian Biol. 31:360-366.  
 376 Eberhardt LS. 1994. Oxygen-consumption during singing by male carolina wrens  
 377 (*Thryothorus ludovicianus*). Auk. 111:124-130.  
 378 Garamszegi LZ, Møller AP. 2004. Extrapair paternity and the evolution of bird song.  
 379 Behav Ecol. 15:508-519.

380 Gaunt AS, Bucher TL, Gaunt SLL, Baptista LF. 1996. Is singing costly? Auk.  
381 113:718-721.

382 Gil D, Gahr M. 2002. The honesty of bird song: multiple constraints for multiple  
383 traits. Trends Ecol Evol. 17:133-141.

384 Gil D, Naguib M, Riebel K, Rutstein A, Gahr M. 2006. Early condition, song learn-  
385 ing, and the volume of song brain nuclei in the zebra finch (*Taeniopygia gut-*  
386 *tata*). J Neurobiol. 66:1602-1612.

387 Grafen A. 1990. Biological signals as handicaps. J Theor Biol. 144:517-546.

388 Hamilton WD, Zuk M. 1982. Heritable true fitness and bright birds - a role for para-  
389 sites. Science. 218:384-387.

390 Houtman AM. 1992. Female Zebra finches choose extra-pair copulations with geneti-  
391 cally attractive males. Proc R Soc Lond B Biol Sci. 249:3-6.

392 Hyman J, Hughes M, Searcy WA, Nowicki S. 2004. Individual variation in the  
393 strength of territory defense in male song sparrows: correlates of age, territory  
394 tenure, and neighbor aggressiveness. Behaviour. 141:15-27.

395 Krebs JR. 1976. Habituation and song repertoires in great tit. Behav Ecol Sociobiol.  
396 1:215-227.

397 Krebs JR, Ashcroft R, Vanorsdol K. 1981. Song matching in the great tit *Parus major*  
398 L. Anim Behav. 29:918-923.

399 Kunc HP, Amrhein V, Naguib M. 2006. Vocal interactions in nightingales, *Luscinia*  
400 *megarhynchos*: more aggressive males have higher pairing success. Anim Be-  
401 hav. 72:25-30.

402 Kwok L. 2003. Felt Tip Sound Studio 2.1. Felt Tip Software.

403 Lambrechts M, Dhondt AA. 1986. Male quality, reproduction, and survival in the  
404 great tit (*Parus major*). Behav Ecol Sociobiol. 19:57-63.

405 Lambrechts M, Dhondt AA. 1988. The anti-exhaustion hypothesis - a new hypothesis  
 406 to explain song performance and song switching in the great tit. *Anim Behav.*  
 407 36:327-334.

408 Langemann U, Tavares JP, Peake TM, McGregor PK. 2000. Response of great tits to  
 409 escalating patterns of playback. *Behaviour.* 137:451-471.

410 McGregor PK, Dabelsteen T, Shepherd M, Pedersen SB. 1992. The signal value of  
 411 matched singing in great tits - evidence from interactive playback experi-  
 412 ments. *Anim Behav.* 43:987-998.

413 McGregor PK, Krebs JR. 1982a. Mating and song types in the great tit. *Nature.*  
 414 297:60-61.

415 McGregor PK, Krebs JR. 1982b. Song types in a population of great tits (*Parus ma-*  
 416 *jor*) - their distribution, abundance and acquisition by individuals. *Behaviour.*  
 417 79:126-152.

418 McGregor PK, Krebs JR, Perrins CM. 1981. Song repertoires and lifetime reproduc-  
 419 tive success in the great tit (*Parus major*). *Am Nat.* 118:149-159.

420 Mennill DJ, Ratcliffe LM. 2004a. Do male black-capped chickadees eavesdrop on  
 421 song contests? A multi-speaker playback experiment. *Behaviour.* 141:125-  
 422 139.

423 Mennill DJ, Ratcliffe LM. 2004b. Overlapping and matching in the song contests of  
 424 black-capped chickadees. *Anim Behav.* 67:441-450.

425 Miyazaki M, Waas JR. 2002. 'Last word' effects of male advertising calls on female  
 426 preference in little blue penguins. *Behaviour.* 139:1413-1423.

427 Naguib M, Fichtel C, Todt D. 1999. Nightingales respond more strongly to vocal  
 428 leaders of simulated dyadic interactions. *Proc R Soc Lond B Biol Sci.*  
 429 266:537-542.

430 Nielsen BMB, Vehrencamp SL. 1995. Responses of song sparrows to song-type  
 431 matching via interactive playback. Behav Ecol Sociobiol. 37:109-117.  
 432 Nowicki S, Hasselquist D, Bensch S, Peters S. 2000. Nestling growth and song reper-  
 433 toire size in great reed warblers: evidence for song learning as an indicator  
 434 mechanism in mate choice. Proc R Soc Lond B Biol Sci. 267:2419-2424.  
 435 Nowicki S, Peters S, Podos J. 1998. Song learning, early nutrition and sexual selec-  
 436 tion in songbirds. Am Zool. 38:179-190.  
 437 Nowicki S, Searcy WA. 2004. Song function and the evolution of female preferences  
 438 - why birds sing, why brains matter. Ann N.Y. Acad Sci. 1016:704-723.  
 439 Nowicki S, Searcy WA, Peters S. 2002. Brain development, song learning and mate  
 440 choice in birds: a review and experimental test of the "nutritional stress hy-  
 441 pothesis". J Comp Physiol A. 188:1003-1014.  
 442 Osiejuk TS, Ratynska K, Cygan JP. 2004. Signal value of alternating and overlapping  
 443 singing in the yellowhammer *Emberiza citrinella*. J Ethol. 22:55-61.  
 444 Osiejuk TS, Ratynska K, Cygan JP. 2007. Corn bunting (*Miliaria calandra*) males  
 445 respond differently to alternating and overlapping playback of song. J Ethol.  
 446 25:159-168.  
 447 Otter K, McGregor PK, Terry AMR, Burford FRL, Peake TM, Dabelsteen T. 1999.  
 448 Do female great tits (*Parus major*) assess males by eavesdropping? A field  
 449 study using interactive song playback. Proc R Soc Lond B Biol Sci. 266:1305-  
 450 1309.  
 451 Peake TM, Matessi G, McGregor PK, Dabelsteen T. 2005. Song type matching, song  
 452 type switching and eavesdropping in male great tits. Anim Behav. 69:1063-  
 453 1068.

454 Richner H, Oppliger A, Christie P. 1993. Effect of an ectoparasite on reproduction in  
 455 great tits. *J Anim Ecol.* 62:703-710.

456 Sall J, Lehmann A. 1996. *JMP Start Statistics*. New York: Duxbury Press.

457 Scheuber H, Jacot A, Brinkhof MWG. 2003a. Condition dependence of a multicom-  
 458 ponent sexual signal in the field cricket *Gryllus campestris*. *Anim Behav.*  
 459 65:721-727.

460 Scheuber H, Jacot A, Brinkhof MWG. 2003b. The effect of past condition on a multi-  
 461 component sexual signal. *Proc R Soc Lond B Biol Sci.* 270:1779-1784.

462 Searcy WA, Andersson M. 1986. Sexual selection and the evolution of song. *Ann Rev*  
 463 *Ecol Syst.* 17:507-533.

464 Spencer KA, Buchanan KL, Goldsmith AR, Catchpole CK. 2003. Song as an honest  
 465 signal of developmental stress in the zebra finch (*Taeniopygia guttata*). *Horm*  
 466 *Behav.* 44:132-139.

467 Spencer KA, Buchanan KL, Goldsmith AR, Catchpole CK. 2004. Developmental  
 468 stress, social rank and song complexity in the european starling (*Stumus vul-*  
 469 *garis*). *Proc R Soc Lond B Biol Sci.* 271:S121-S123.

470 Spencer KA, Buchanan KL, Leitner S, Goldsmith AR, Catchpole CK. 2005a. Para-  
 471 sites affect song complexity and neural development in a songbird. *Proc R Soc*  
 472 *Lond B Biol Sci.* 272:2037-2043.

473 Spencer KA, Wimpenny JH, Buchanan KL, Lovell PG, Goldsmith AR, Catchpole  
 474 CK. 2005b. Developmental stress affects the attractiveness of male song and  
 475 female choice in the zebra finch (*Taeniopygia guttata*). *Behav Ecol Sociobiol.*  
 476 58:423-428.

477 Todt D. 1981. On functions of vocal matching - effect of counter-replies on song post  
 478 choice and singing. *Z Tierpsychol.* 57:73-93.

479 Todt D, Naguib M. 2000. Vocal interactions in birds: The use of song as a model in  
 480 communication. In: Advances in the Study of Behavior, Vol. 29. 247-296.  
 481 Tripet F, Richner H. 1997a. Host response to ectoparasites: Food compensation by  
 482 parent blue tits. Oikos. 78:557-561.  
 483 Tripet F, Richner H. 1997b. The coevolutionary potential of a 'generalist' parasite, the  
 484 hen flea *Ceratophyllus gallinae*. Parasitology. 115:419-427.  
 485 Vehrencamp SL. 2001. Is song-type matching a conventional signal of aggressive  
 486 intentions? Proc R Soc Lond B Biol Sci. 268:1637-1642.  
 487 Vehrencamp SL, Bradbury JW, Gibson RM. 1989. The energetic cost of display in  
 488 male sage grouse. Anim Behav. 38:885-896.  
 489

## FIGURE LEGENDS

Figure 1 Spectrogram of a typical great tit song.

Figure 2 Relationship between song duration and song overlap of adult males raised in a flea-infested ( $n = 13$ ; black) or a parasite-free nest ( $n = 9$ ; white). The triangles indicate old males (year of birth shown), the points indicate one-year old males (born in 2004). Song overlap (% number of strophes overlapping with playback) and song duration were recorded during a 20-minute period in response to a playback.

Figure 3 Relationship between song duration and clutch size. Black signs indicate males infested with fleas early in life ( $n = 13$ ), white signs represent uninfested males ( $n = 9$ ). The triangles indicate old males (year of birth shown), the points indicate one-year old males (born in 2004). Song duration was recorded during a 20-minute period in response to a playback.

**FIGURES**

Figure 1






Figure 2





514    Figure 3



515

516